ASD, Parental Grief and Adaptation

ASD diagnosis and Parental grief

When parents learn their child has autism, they often experience a deep sense of loss—fearing their child may never develop as they hoped. This reaction is not just sadness, but a powerful surge of emotions: fear, disbelief, and grief. The grief parents feel is often compared to mourning the loss of a loved one, echoing the process outlined in Attachment Theory, which explains how strong emotional bonds lead to intense grief when disrupted (Bretherton, 1992).

Loss of expectations and Grief

The bonds parents form with their children are rooted in security and safety; they are not just emotional—they're biological. When parents lose their expectations about their child's future, their reactions, crying, anger, yearning, are survival mechanisms, meant to restore the lost connection (Bowlby, 1980). Here is what a mother might typically experience when first finding out:

The Adaptation Process

Adapting to a diagnosis is a journey, not a single event. Four main tasks define this process:

1. **Accepting the reality:** Parents face and accept that their child is neurologically different. Denial is common, some may downplay the diagnosis or its impact, delaying assessment and therapy.
2. **Processing emotional pain:** Allowing oneself to feel sadness, anger, or loss helps, though some communities discourage open grieving.

3. **Adjusting to new roles:** Parents may have to develop new skills and routines, which can challenge their self-image and sense of competence.
4. **Emotionally relocating:** Parents need to form a new, ongoing relationship with their child, while still moving forward with their own lives (Worden, 2009).

Complicated Adaptation

Sometimes, the grieving process stalls or becomes overwhelming, leading to chronic, delayed, exaggerated, or masked grief. This may be triggered by multiple simultaneous losses, personal history, personality factors, or lack of support and resources (Shear, 2015). Therapy focusses on resolving these stuck points, sometimes by revisiting earlier or unresolved losses.

Healthy Adaptation

Adaptation is considered healthy when parents can find pleasure again, take on new roles, and think about their child without overwhelming pain, though moments of sadness may still arise (Worden, 2009).

How Psychologists can help

Psychologists can play a crucial role in supporting parents and families after a child is diagnosed with autism. Here's how they can help, with research-backed approaches and references:

1. Provide Psychoeducation

Psychologists can help parents understand what autism is, what it isn't, and what to expect. They help dispel myths, explain the range of possible outcomes, and help families set realistic expectations (Karst & Van Hecke, 2012). This reduces fear and uncertainty, and empowers parents to make informed decisions.

2. Offer Emotional Support and Grief Counselling

Receiving an autism diagnosis can trigger grief, anxiety, guilt, anger, and depression. Psychologists can provide a non-judgmental space for parents to process these emotions, using evidence-based grief counselling techniques and teaching coping strategies that help normalise these feelings.

3. Strengthen Coping Skills

Therapists teach parents skills for managing stress and emotional overload. This might include mindfulness, cognitive-behavioral tools for reframing negative thoughts, and relaxation exercises.

4. Facilitate Family Communication and Adjustment

A diagnosis can strain marriages and family relationships. Psychologists can work with the whole family to improve communication, resolve conflicts, navigate new roles, and foster resilience.

5. Support Behaviour Management and Parenting Strategies

Psychologists can teach parents positive behaviour support strategies, help them respond to challenging behaviours, and set up effective routines at home. They can also coach parents in using evidence-based interventions tailored to their child's needs.

6. Identify and Treat Mental Health Issues

If a parent develops depression, anxiety, or experiences prolonged grief, psychologists can provide individual therapy or refer to a psychiatrist if needed. Early intervention prevents long-term problems.

7. Advocate for the Family

If needed, psychologists can help advocate for the child and family within schools, healthcare systems, and the broader community, ensuring that the child's needs are met and that parents' voices are heard.

(03) 9848 9100

Updated on: 19/02/2026 by: Dr. Jacques Duff PhD – BA Psych; Grad Dip Psych; MECNS; MAAAPB; MISNR; FANSA

Reviewed on: 23/02/2026 by: Bernard Ferriere - Clinical Psychologist; BA; Grad Dip App Psych; Dip Clinical Hypnosis; MAPS;