Can Psychologists Diagnose Autism in Australia?



A parent may have noticed that their child has found social situations difficult since preschool. An adult may recognise lifelong patterns of sensory overload, intense interests or exhaustion after masking at work. In either situation, a practical question often comes first: **can psychologists diagnose autism?**

In Australia, a psychologist may diagnose autism when they have the appropriate training, experience and clinical scope to conduct a comprehensive assessment. Autism assessment is not based on a single questionnaire, a short appointment or an

online checklist. It involves careful consideration of a person's developmental history, current presentation, strengths, support needs and other possible explanations for their experiences.

For some people, a psychologist-led assessment is appropriate. For others, the clearest pathway involves input from a paediatrician, psychiatrist, speech pathologist, occupational therapist or other relevant professionals. The right arrangement depends on the person's age, presentation, medical history and the purpose of the assessment.

Can psychologists diagnose autism independently?

Psychologists are trained to assess behaviour, cognition, development, emotional wellbeing and mental health. Registered psychologists who work within their competence in autism assessment can identify whether a person meets the recognised diagnostic criteria for autism spectrum disorder.

A diagnosis should be based on a thorough clinical process, usually guided by the current Diagnostic and Statistical Manual of Mental Disorders criteria. The psychologist considers whether social communication differences and restricted, repetitive patterns of behaviour, interests or sensory responses have been present from early development, cause meaningful functional impact, and are better explained by another condition alone.

There is no requirement that every autism assessment be completed by a medical practitioner. However, medical and multidisciplinary input can be valuable, and may be necessary in particular circumstances. For example, a paediatrician or psychiatrist may assess co-occurring medical, developmental or mental health concerns, consider medication where relevant, or provide documentation required by a particular service.

It is reasonable to ask a clinician about their experience with autism assessment in the relevant age group. Adult autism presentations can differ from those recognised in childhood, particularly for people who have learned to mask or compensate

socially. Similarly, children with strong language skills or high academic ability may still have support needs that deserve careful assessment.

What a comprehensive autism assessment involves

A considered assessment gathers information from more than one source. The process is tailored to the individual, but it commonly includes a detailed clinical interview, developmental history and standardised assessment measures where appropriate.

For a child, parents or caregivers are usually central to the developmental history. The psychologist may also seek information from school, with consent, because classroom demands can reveal differences that are not as apparent at home. Observations, reports from teachers and information about learning, friendships, play, behaviour and sensory experiences can all contribute to a more complete picture.

For adults, assessment often involves exploring childhood development as well as current life. A partner, parent or sibling may be invited to contribute information if the adult is comfortable with this and someone is available. School reports, earlier assessments or other records can be helpful, but their absence does not automatically prevent an assessment. Many adults seeking clarity were not identified when they were at school.

The psychologist will also consider factors that may overlap with, accompany or sometimes resemble autism. These can include ADHD, anxiety, depression, trauma, language differences, learning difficulties, intellectual disability, obsessive-compulsive symptoms and sensory processing concerns. This is not about looking for reasons to dismiss a person’s experience. It is about reaching an explanation that is clinically sound and useful for planning support.

Standardised tools can support assessment, but they do not replace professional judgement. A score on a screening questionnaire cannot diagnose autism by itself. Equally, a person may have a presentation that warrants detailed consideration even

if a brief screener does not clearly indicate autism.

Why assessment pathways can look different

Autism is a spectrum, but that phrase does not mean every person has the same set of traits or requires the same kind of support. Some people need substantial assistance with daily living, communication or emotional regulation. Others may manage work, study and relationships while experiencing considerable effort, fatigue or misunderstanding that is not visible to others.

Age is one reason pathways differ. Young children may need developmental assessment that considers speech, play, adaptive skills and family observations. Adolescents may need an assessment that accounts for school pressures, peer relationships, anxiety and identity development. Adults may be seeking an explanation for longstanding patterns that have previously been attributed to shyness, anxiety, ADHD or personality.

The intended use of the report also matters. A report prepared to guide therapy, school adjustments or workplace discussions may need different detail from one being considered for disability support access. Requirements can change between organisations and funding systems, so it is sensible to clarify what documentation is needed before commencing an assessment.

Psychologist, paediatrician or psychiatrist: who should you see?

There is no single answer for every family or adult. A psychologist may be well placed to undertake a detailed autism assessment, particularly where the main need is diagnostic clarification and recommendations for psychological, behavioural or practical supports.

A paediatrician may be involved when a child has broader developmental, medical or physical health questions that require medical assessment. A psychiatrist may be particularly relevant where there are complex mental health concerns, diagnostic uncertainty, or a need to consider medication for a co-occurring condition. Neither

paediatricians nor psychiatrists automatically replace the detailed psychological assessment that can be needed to understand a person's day-to-day functioning.

A multidisciplinary assessment can be appropriate when communication, motor, sensory, learning or adaptive functioning questions need specialised input. While it can provide a broad perspective, it may also involve additional appointments and cost. A clinician should be able to explain why particular professional input is recommended rather than assuming every person needs the same process.

What happens after an autism diagnosis?

A diagnosis is not the end point of care. It should provide a clearer framework for understanding needs, making informed choices and accessing appropriate support. For some people, the most useful next step is psychological therapy focused on anxiety, emotional regulation, self-understanding or coping with change. For children, parents may benefit from practical strategies that support communication, routines, school participation and behaviour without trying to make the child appear less autistic.

Recommendations should be individualised. They may address school adjustments, workplace communication, sensory needs, executive functioning, relationships, sleep, allied health support or management of co-occurring ADHD and anxiety. Not every recommendation will be necessary, and support plans can change over time.

At Behavioural Neurotherapy Clinic, assessment and therapy can be considered as part of an ongoing clinical pathway rather than as a diagnosis-only process. Appointments are available in Melbourne and, where clinically appropriate, through Australia-wide telehealth consultations. The suitability of telehealth depends on the person's age, assessment needs and available information.

Referrals, Medicare and preparing for an appointment

You do not always need a referral to see a psychologist privately, although a referral from a GP, paediatrician, psychiatrist or another health professional can provide

useful background. Medicare rebates may be available for eligible psychological services when a person has an appropriate referral and Mental Health Care Plan. The specific rebate pathway and whether it applies to assessment work should be confirmed before booking, as arrangements vary.

Before an appointment, it can help to gather relevant information without feeling pressure to produce a perfect file. School reports, previous assessments, medical letters and a brief list of examples from everyday life may assist. For adults, reflecting on childhood friendships, sensory experiences, routines, school and work patterns can provide useful starting points. For parents, noting what happens before, during and after challenging situations may help describe a child's needs clearly.

Seeking an assessment does not presume a diagnosis. It creates space for a careful conversation about what is happening, what information is needed and what support may be useful. A well-explained assessment process can offer a more grounded next step than trying to make sense of complex experiences alone.

For more information

Call (03) 9848 9100

Published: 28/08/2026 and updated on: 17/10/2026 by Dr Jacques Duff-PhD; BA Psych; Grad Dip Applied Psychology; FANSA; Clinical Neuroscientist