Can Anxiety Mimic ADHD? The Key Differences



A child who cannot settle to homework, loses track of instructions and becomes upset before school may appear to have ADHD. An adult who procrastinates, misses details in meetings and feels constantly overwhelmed may wonder the same thing. So, can anxiety mimic ADHD? In some situations, yes. Anxiety can affect concentration, memory, organisation and emotional regulation in ways that resemble ADHD. The reverse can also happen: the daily demands of living with ADHD can contribute to anxiety.

The overlap can be confusing, particularly when someone has spent years trying to understand why tasks that seem straightforward for others feel disproportionately difficult. A careful assessment looks beyond a single symptom and considers when difficulties began, where they occur, what triggers them and how they have changed over time.

Can anxiety mimic ADHD in children and adults?

Anxiety is more than occasional worry. It can involve persistent fears, racing thoughts, physical tension, avoidance, reassurance-seeking and a sense of being on alert. When the mind is preoccupied with possible problems, there is less attention available for the task at hand.

This may look like distractibility. A student may stare out the window because they are worrying about making a mistake. An adult may repeatedly reread an email because they fear getting it wrong, then run out of time for other work. Someone who is anxious may also seem forgetful because information was not fully taken in while they were tense or mentally preoccupied.

Sleep disruption is another common point of overlap. Worry can make it harder to fall asleep or return to sleep after waking. Tiredness can then reduce concentration, patience, planning and impulse control the next day. In children, anxiety may also show up as irritability, restlessness, school refusal, physical complaints or difficulty separating from a parent or caregiver.

These experiences are real and can significantly affect daily life. However, symptoms that resemble ADHD do not automatically indicate ADHD, anxiety or any other particular condition. Context and developmental history matter.

Where the symptoms overlap

Both anxiety and ADHD can be associated with trouble concentrating, restlessness, forgetfulness, interrupted sleep, irritability and difficulty completing tasks. Emotional reactions can also be misunderstood. A person who becomes distressed by an unexpected change may be responding to anxious anticipation, difficulty shifting

attention, sensory overload, a previous negative experience, or more than one of these factors.

Executive functioning difficulties are especially relevant. Executive functions include skills such as starting tasks, organising materials, estimating time, holding information in mind and regulating responses. ADHD is commonly associated with ongoing difficulties in these areas. Anxiety may interfere with the same skills when worry, perfectionism or avoidance takes up mental energy.

The outward behaviour may therefore be similar, while the underlying driver differs. For example, delayed work may stem from fear of failure and a need for certainty. It may stem from longstanding difficulty initiating, sequencing and sustaining attention. It may also reflect both. This distinction guides appropriate support, so it is not something to resolve through a symptom checklist alone.

Patterns that may point more strongly towards anxiety

With anxiety, concentration difficulties often become more pronounced around particular worries or situations. These may include school performance, social judgement, health concerns, separation, conflict or uncertainty. A person may focus well when they feel safe, calm and interested, but struggle sharply when a feared event is approaching.

Avoidance can be a significant feature. The individual may delay a task because beginning it brings up worry, seek repeated reassurance, or avoid situations where they believe they might be judged or make an error. Physical symptoms such as a racing heart, nausea, muscle tension or stomach pain can also provide useful clinical context, although they are not present for everyone.

Patterns that may point more strongly towards ADHD

ADHD is a neurodevelopmental condition, which means relevant traits are generally present from childhood, even if they were not recognised at the time. In adults seeking assessment, the pattern may become clearer when they reflect on early school reports, family recollections, unfinished tasks, disorganisation or feedback about being distracted, impulsive or excessively talkative.

ADHD-related attention and executive-function difficulties are usually evident across more than one setting, such as home, school, university, work or daily administration. They may persist even when a person is doing something they value and is not especially worried. At the same time, attention can be highly variable. Some people can concentrate intensely on an engaging activity yet struggle to begin routine, repetitive or less rewarding tasks.

This is not a simple rule. Anxiety can occur across many settings, and people with ADHD can become particularly anxious in situations where past disorganisation, lateness or criticism has had consequences. A qualified clinician considers the overall pattern rather than relying on one apparent difference.

Why ADHD and anxiety can occur together

It is possible to have ADHD and an anxiety disorder at the same time. For some people, anxiety develops in part from repeated experiences of falling behind, forgetting obligations, misreading time or feeling unable to keep up with everyday expectations. For others, anxiety is a separate difficulty that needs its own assessment and treatment plan.

Co-occurring concerns can affect how symptoms present. A highly anxious child with ADHD may appear quiet and compliant at school while struggling at home after holding everything together during the day. An adult may rely on worry and last-minute urgency to start tasks, which can mask underlying attention or planning difficulties for years. Conversely, someone with anxiety may over-prepare and develop rigid systems that make their concentration difficulties less visible.

A comprehensive assessment should also consider mood, sleep, trauma history, learning differences, autism, substance use, physical health factors and medication effects where relevant. The aim is not to attach as many labels as possible. It is to develop a clinically sound understanding of the factors affecting functioning.

What a comprehensive ADHD assessment considers

A formal assessment is more detailed than an online screener or a short discussion of current symptoms. Screening tools can be useful starting points, but they cannot determine the cause of concentration difficulties on their own.

Assessment commonly includes a detailed clinical interview, developmental and family history, discussion of current functioning, standardised questionnaires and information from other settings where available. For children and adolescents, this may include parent, caregiver and school input. For adults, relevant information may include early educational experiences, reports where available and the impact of symptoms on work, relationships, study, finances and daily responsibilities.

Clinicians also explore the timing of symptoms. Did attention difficulties occur before anxiety became prominent? Are they present during calmer periods? Does a person avoid tasks because they cannot organise a starting point, because they fear the outcome, or because both processes are operating? These questions help distinguish overlapping presentations without reducing a person's experience to a single explanation.

Choosing practical next steps

If concentration or behaviour changes suddenly, is accompanied by marked changes in mood, sleep or physical health, or is creating immediate safety concerns, timely medical advice is appropriate. For ongoing difficulties, discussing concerns with a GP can help determine whether further psychological, psychiatric, paediatric or other health assessment is indicated.

For people considering an ADHD assessment, it can be useful to bring examples rather than trying to prove a diagnosis. Note situations where attention, organisation, restlessness, worry or emotional regulation create difficulty. Parents may wish to record observations from home and school. Adults can reflect on patterns across childhood and adult life, including what strategies they use to manage demands and the effort those strategies require.

At Behavioural Neurotherapy Clinic, comprehensive assessment and therapy can be accessed in Melbourne or through Australia-wide Zoom consultations. Eligible people

may be able to access Medicare rebates with an appropriate GP, psychiatrist or paediatrician referral and Mental Health Care Plan.

Clarity often begins with recognising that distractibility is not a character flaw, and that similar-looking symptoms can have different causes. A thoughtful assessment can make room for the full picture, then support decisions about care that fit the individual rather than the label alone.

For more information

Call (03) 9848 9100

Published: 07/09/2026 and updated on: 23/11/2026 by Dr Jacques Duff-PhD; BA Psych; Grad Dip Applied Psychology; FANSA; Clinical Neuroscientist