ADHD Versus Anxiety Symptoms: How They Differ



A child who cannot start homework may be distracted by every sound in the room, worried about getting the answers wrong, or both. An adult who misses deadlines may be struggling to organise tasks, avoiding work that feels threatening, or managing years of accumulated stress. **ADHD versus anxiety symptoms** can be difficult to separate because both may affect concentration, sleep, emotional regulation and daily functioning.

The distinction matters because attention-deficit/hyperactivity disorder (ADHD) and anxiety can require different clinical approaches. They can also occur together. A careful assessment looks beyond a checklist of symptoms to understand when difficulties began, what triggers them, how they appear across settings, and the effect they have on school, work, relationships and everyday life.

Why ADHD and anxiety can look alike

Both ADHD and anxiety can make it hard to focus. A person may appear restless, forget appointments, lose track of conversations, procrastinate, or become overwhelmed by tasks that seem manageable to others. Children may be described as distracted, oppositional or overly sensitive. Adults may feel they are always behind, despite considerable effort.

The outward behaviour is not always a reliable guide to the underlying reason. For example, a student who looks away during class could be following unrelated thoughts associated with ADHD, worrying about being called on, struggling with a learning difficulty, not sleeping well, or experiencing a combination of these factors.

Emotional responses can overlap too. Frustration after making a mistake, irritability when routines change and difficulty settling at night may occur in either condition. This is why online questionnaires and isolated symptom lists can be useful prompts for reflection, but cannot establish a diagnosis.

ADHD versus anxiety symptoms: key differences

ADHD is a neurodevelopmental condition. Its core features relate to patterns of inattention and/or hyperactivity-impulsivity that begin in childhood, even if they are only recognised later. These patterns are generally evident in more than one area of life and affect functioning over time.

Anxiety involves excessive fear, worry or apprehension that can be difficult to control. It may be focused on particular situations, such as social interaction or separation, or take the form of more persistent and wide-ranging worry. Physical sensations such as a racing heart, muscle tension, nausea or feeling on edge can be

prominent, although they are not present for everyone.

What attention difficulties may feel like in ADHD

With ADHD, attention may shift quickly even when the person is not especially worried. There can be difficulty getting started, sequencing steps, estimating time, remembering instructions and maintaining effort on tasks that are repetitive or not immediately engaging. Some people can concentrate deeply on activities that strongly interest them, then find routine demands unusually hard to begin or complete.

Impulsivity may show up as interrupting, acting before considering consequences, making quick decisions, or speaking more than intended. Hyperactivity is not limited to visibly high activity levels. In adolescents and adults, it can feel like internal restlessness, a need to keep busy, or difficulty relaxing.

The pattern is often longstanding. A parent may recall early concerns about distractibility, activity level, forgetfulness or emotional reactivity. An adult seeking later-life assessment may recognise similar difficulties across school years, work roles and home responsibilities, even if they found ways to compensate.

What concentration difficulties may feel like in anxiety

With anxiety, concentration is often disrupted because attention is pulled towards threat, uncertainty or anticipated negative outcomes. A person may repeatedly review conversations, imagine what could go wrong, seek reassurance, avoid decisions or delay tasks because they fear making an error.

Avoidance is often a useful clinical clue. Someone may be capable of completing a task but feel unable to approach it when worry is high. Once the source of anxiety reduces, their concentration and organisation may improve substantially. That does not rule out ADHD, but it helps clarify the role anxiety may be playing.

Anxiety can also be episodic. Symptoms may increase around exams, workplace conflict, health concerns, major changes or social demands. ADHD-related executive functioning difficulties are more likely to be present across different circumstances,

although stress, sleep and workload can make them more noticeable.

When ADHD and anxiety occur together

Having one condition does not protect someone from the other. Some people with ADHD develop anxiety after repeated experiences of missed deadlines, criticism, academic difficulty, social misunderstandings or the effort of trying to keep up with daily demands. Others have anxiety that makes ADHD traits more difficult to manage.

When both are present, symptoms can pull in different directions. ADHD-related impulsivity may lead a person to act quickly, while anxiety may make them overthink the same decision afterwards. A child might want to join an activity but struggle with the planning and transition involved, then appear reluctant or avoidant. An adult may postpone an important task due to worry and also find it difficult to organise the steps needed to complete it.

This overlap means that treatment planning needs to be individualised. Addressing anxiety without considering executive functioning difficulties may leave practical barriers unresolved. Equally, focusing only on attention and organisation may overlook persistent worry, avoidance or physical anxiety symptoms.

What a comprehensive assessment considers

A clinical assessment is not based on one appointment, one school report or a single screening tool. The process usually brings together developmental history, current concerns, functional impact and information from relevant settings. For children and adolescents, this may include parent input and school-based information where appropriate. For adults, it may include a detailed history of childhood experiences, education, employment, relationships and current responsibilities.

Clinicians also consider other explanations for overlapping symptoms. Sleep difficulties, depression, trauma experiences, autism, learning differences, substance use, medical conditions and significant life stress can affect attention, mood and behaviour. The goal is not to assign every difficulty to one label. It is to develop a

clinically sound understanding of what is contributing and what support may be appropriate.

A formal ADHD assessment generally considers whether symptoms were present in childhood, occur across settings and cause meaningful impairment. Anxiety assessment considers the nature of fears and worries, physical symptoms, avoidance patterns and how long these have been present. Where symptoms are complex, assessment may take more than one consultation.

Signs it may be useful to seek professional guidance

It may be worth discussing concerns with a GP or qualified mental health professional when attention, worry, behaviour or emotional distress is persistently affecting learning, work, relationships or family life. For a child, this could include ongoing difficulty managing school demands, frequent meltdowns around transitions, avoidance of activities they would otherwise enjoy, or a marked change in confidence. For an adult, it may involve chronic disorganisation, escalating worry, difficulty maintaining routines, or feeling exhausted by the effort required to manage ordinary tasks.

The timing and context are important. Sudden changes in concentration or behaviour warrant consideration of current stressors, health and sleep, rather than an assumption that ADHD is the cause. Conversely, a person who has always struggled but has reached a point where their usual strategies no longer work may benefit from a structured assessment.

For people in Melbourne, in-person assessment may be one option; telehealth can also support access for eligible clients elsewhere in Australia, depending on clinical needs and appointment suitability. A GP or specialist can advise on referral pathways. Medicare rebates may be available for eligible psychological services when a person has an appropriate referral and Mental Health Care Plan.

Support starts with a clearer picture

Support for ADHD may include psychological strategies for planning, organisation, routines, emotional regulation and behaviour, alongside medical review where indicated. Support for anxiety commonly includes psychological therapy that helps a person understand worry patterns, reduce avoidance and build practical coping skills. When conditions co-occur, treatment can be coordinated around the difficulties causing the greatest impact first, while still recognising the full picture.

For parents, it can help to describe specific observations rather than trying to decide which diagnosis fits: when the difficulty occurs, what happens beforehand, how long it has been present and what helps. Adults can do the same by noting patterns across work, home and earlier life. Clear information gives a clinician a stronger starting point than self-diagnosis alone.

Feeling uncertain about whether symptoms reflect ADHD, anxiety or both is common. A thoughtful assessment can replace guesswork with an informed understanding and a practical path for next steps.

For more information

Call (03) 9848 9100

Published: 20/09/2026 and updated on: 17/10/2026 by Dr Jacques Duff-PhD; BA Psych; Grad Dip Applied Psychology; FANSA; Clinical Neuroscientist