ADHD Assessment Examples: What Clinicians Look For



A school report that describes unfinished work, a parent's notes about difficult mornings, and an adult's memories of always losing track of tasks can all be useful ADHD assessment examples. None of these items provides a diagnosis on its own. Together with structured measures, developmental history and clinical interviews, they help a psychologist understand whether ADHD may explain a pattern of difficulties, or whether another factor needs attention.

For parents and caregivers, the process can feel unfamiliar after a school raises concerns or daily routines become harder to manage. A child ADHD assessment is not a test of parenting, effort or character. It is a structured process for understanding how attention, activity level, impulsivity, learning, emotions and everyday functioning fit together over time.

Why one example is never enough

ADHD can look different across children and settings. A child may be settled during a preferred activity but struggle to start written work. Another may appear highly active at home yet become quiet and overwhelmed at school. Some children work very hard to keep up during the day and show their fatigue or frustration later.

For this reason, clinicians look for a persistent pattern rather than relying on a single behaviour. They consider whether difficulties were present during development, occur in more than one setting, and interfere with daily functioning. The assessment also considers other explanations that can overlap with ADHD-like difficulties, including anxiety, sleep problems, learning difficulties, autism, hearing or vision concerns, changes at school or home, and emotional distress.

A behaviour such as interrupting may have several possible meanings. It can relate to impulsivity, but it may also occur when a child is anxious, excited, has difficulty reading social timing, or is trying to hold onto a thought before it disappears. The context matters.

ADHD assessment examples from everyday life

The examples below show the kind of information a psychologist may explore. They are not checklists for diagnosing ADHD at home, and no individual example confirms or rules out a diagnosis.

Attention and task completion

A parent or teacher may describe work that begins well but is left unfinished, especially when several steps are involved. The child might need repeated reminders

to pack a bag, complete a homework routine or return to a task after an interruption. In class, a teacher may note missed instructions, avoidable errors, slow completion, or a tendency to look around the room when independent work begins.

The clinician will ask for detail. It helps to know whether the difficulty occurs only with certain subjects, after poor sleep, in noisy environments, or across tasks that the child enjoys as well as tasks they dislike. This distinction can help separate attention regulation difficulties from a specific learning difficulty or a mismatch between task demands and current skills.

Organisation and executive functioning

Executive functioning refers to the mental skills used to plan, begin, remember and finish tasks. Assessment examples may include a bedroom or school bag that becomes disorganised quickly, regular loss of lunch boxes or sports gear, and difficulty estimating how long a task will take.

At school, this might show up as incomplete projects despite good ideas, trouble following a timetable, or forgetting materials needed for the next lesson. At home, it may appear as a routine that works only with close adult support. Clinicians consider the level of assistance required, not simply whether the task is eventually completed.

Activity level and impulse control

Some children move frequently, talk a great deal, call out, climb, fidget or find waiting especially difficult. Others show less obvious restlessness. They may doodle constantly, tap, shift position, rush through work, or seem unable to settle their thoughts at bedtime.

Examples involving impulsivity can include acting before hearing the full instruction, moving quickly from an idea to an action, or becoming upset after a consequence that seemed unexpected to the child. A clinician considers whether these behaviours are developmentally unusual, how often they occur, and whether they create difficulties with learning, relationships or safety.

Emotional regulation and recovery after stress

Emotional regulation is commonly explored because some children have strong reactions to frustration, changes in plans, criticism or small mistakes. An assessment may examine the build-up to these moments, how long recovery takes, and what support helps.

This does not mean that emotional reactions automatically indicate ADHD. Anxiety, autism, learning demands, family stress, sensory sensitivities and sleep difficulties can all affect regulation. A careful assessment keeps these possibilities open rather than fitting every difficulty into one explanation.

Social and family observations

Parents and caregivers often notice patterns that school staff do not see, particularly during mornings, homework, meals, transitions and bedtime. Teachers may have a different but equally useful view of concentration, peer interactions, classroom expectations and academic progress.

When appropriate and with consent, information from both settings can help show whether a pattern is widespread or mainly linked to one environment. A psychologist may also ask about friendships, conflict, play, family routines and the strategies already being used. The purpose is to understand functioning, not to judge the family.

What a child ADHD assessment usually includes

The exact process depends on the child's age, presentation and available information. It commonly begins with an appointment for parents or caregivers to discuss developmental history, current concerns, health background, family context and school experience. Developmental history may cover early milestones, language, sleep, behaviour, learning and previous supports.

The psychologist may then meet with the child or adolescent. This appointment can include conversation, observation and activities selected to understand attention, thinking skills, learning or emotional wellbeing. Structured questionnaires may be completed by parents, teachers and, for older children, the young person. School

reports, work samples and previous assessments can also provide useful context.

A cognitive or learning assessment may be considered when there are concerns about reading, writing, mathematics, working memory or processing speed. This is particularly relevant when a child's effort and apparent ability do not match their school progress. An autism assessment or assessment of anxiety and mood may also be discussed when the developmental history and current presentation indicate this would be useful.

The assessment is usually completed over more than one appointment. A feedback session explains the information gathered, the clinical formulation and any diagnostic support offered. Families can ask about the practical meaning of the findings for home, school and wellbeing.

What happens when ADHD is considered alongside other factors

A clear result is not always possible from the first set of information. For example, significant sleep disruption can affect concentration and emotional regulation. A child who has fallen behind in reading may avoid written work because the task is genuinely difficult. Anxiety can make a child appear distracted, restless or unable to begin.

In these situations, the clinician may recommend further assessment, school information, psychological support or discussion with another health professional. This is not a failed assessment. It reflects the fact that children's development and daily functioning are influenced by more than one factor.

Where diagnostic support for ADHD is provided, planning can include behavioural strategies, parent support, school adjustments, psychological therapy for related anxiety or emotional difficulties, and coordination with the child's GP, paediatrician or psychiatrist. Psychologists at BNC provide assessment, diagnostic support and psychological therapy. Medical decisions, including whether a prescriber considers medication, sit with the relevant medical practitioner.

Preparing useful information for an appointment

Families do not need to create a perfect record before seeking an assessment. Ordinary documents and observations are often the most useful. Recent and past school reports, teacher comments, work samples, details of previous assessments, and notes about developmental or health history can help build a fuller picture.

Brief examples are generally more helpful than broad labels. For instance, noting that a child needed an adult beside them for each step of a homework task gives more clinical information than saying they are “lazy” or “unmotivated”. Recording what happened before a difficulty, what the child did, and what helped afterwards can also give the clinician a clearer starting point.

For adolescents, it can be helpful to include their own account of school, friendships, routines and goals. Their perspective may differ from the adults around them, and both views matter.

A pathway built around the whole picture

ADHD assessment examples are useful because they turn a general concern into information that can be understood in context. They do not need to be dramatic. Repeated small difficulties with starting, remembering, organising, regulating or recovering can be meaningful when considered alongside developmental history and daily functioning.

A well-explained assessment gives families language for what has been happening and a practical basis for deciding what support may be appropriate next. Whether ADHD is confirmed, not confirmed, or considered alongside other needs, the next step should be guided by the child’s full picture rather than a single label.

For more information

Call (03) 9848 9100

Published: 08/10/2026 and updated on: 08/10/2026 by Dr Jacques Duff-PhD; BA Psych; Grad Dip Applied Psychology; FANSA; Clinical Neuroscientist