Does QEEG Diagnose ADHD? What Assessment Needs



A brain map can look precise: coloured images, graphs and a report comparing patterns of electrical activity with a reference database. For families seeking clarity, it may seem like a direct answer. But does QEEG diagnose ADHD? No. A quantitative electroencephalogram, or QEEG, may provide additional information in selected circumstances, but it cannot diagnose ADHD on its own.

ADHD is identified through a clinical assessment of a person’s developmental history, current functioning and the pattern of difficulties across settings. There is no single

brain scan, blood test, questionnaire or computer task that can confirm or rule out ADHD. A meaningful assessment considers the whole picture, including other explanations that may need attention.

What a QEEG measures

An electroencephalogram, commonly called an EEG, records the brain's electrical activity through sensors placed on the scalp. A QEEG uses computer analysis to compare features of that recording, such as activity across different frequency bands, with a reference database. It is sometimes described as brain mapping.

The output may show that aspects of a recording differ from the reference group used by the system. That finding is not the same as identifying ADHD. Brain activity varies between people and can also be affected by sleep, fatigue, anxiety, medication, caffeine, age, the recording conditions and technical factors.

Importantly, patterns sometimes associated with ADHD are not unique to ADHD. Similar findings may occur in people with other conditions or in people without a diagnosis. A QEEG also cannot establish whether difficulties began in childhood, occur across home and school or work, or create the functional impact required for an ADHD diagnosis.

Does QEEG diagnose ADHD in Australia?

QEEG should not be used as a stand-alone diagnostic test for ADHD. A diagnosis requires clinical judgement by an appropriately qualified clinician using recognised diagnostic criteria. The assessment needs to establish a persistent pattern of inattention and/or hyperactivity-impulsivity, the age at which relevant difficulties began, the settings in which they occur, and their impact on daily life.

This matters because attention and executive functioning can be affected by many factors. Insufficient sleep, anxiety, depression, trauma, learning difficulties, autism, hearing or vision concerns, medical conditions, substance use and major life stress can all influence concentration, organisation, memory or emotional regulation. These factors may occur alongside ADHD as well.

A brain map cannot sort through that context. It cannot hear a parent describe early development, consider school reports over time, or explore why an adult's difficulties became more noticeable when work, study, parenting or household demands increased. Those details are central to diagnostic support.

What an ADHD assessment involves instead

For a child or adolescent, assessment commonly includes information from parents or caregivers, developmental history and structured rating scales. School information can be useful because ADHD-related difficulties need to be considered across more than one setting. The psychologist also considers learning, social communication, behaviour, emotional wellbeing, sleep and family circumstances where relevant.

For adults, the process similarly looks beyond a current list of symptoms. It includes childhood history, education and work experiences, relationships, daily organisation, emotional regulation and the practical effects of the reported difficulties. When available and appropriate, input from someone who knew the person as a child may add useful context. It is understandable if this is not possible for every adult seeking assessment.

Structured interviews and standardised measures help organise this information. They do not replace clinical judgement, but they provide a consistent way to examine whether the reported pattern aligns with diagnostic criteria. A psychologist may also recommend assessment for learning difficulties, autism or mental health concerns when the history indicates that these areas need consideration.

The aim is not to fit a person into a label quickly. It is to understand what is contributing to the difficulties and to provide clear documentation and practical next steps.

When QEEG may be discussed

A QEEG may be raised by a provider offering neurofeedback or by a family who has encountered brain mapping online. It may also be discussed when there is a separate clinical reason to investigate neurological symptoms. Those are different purposes

from ADHD diagnosis.

If QEEG is being considered, it is reasonable to ask what question the recording is intended to answer, who will interpret it, and how its findings will be used alongside a full clinical assessment. It is also reasonable to ask whether the provider is presenting QEEG as one source of information or as a diagnostic shortcut.

Families and adults should be cautious about claims that a particular QEEG pattern proves ADHD, identifies a subtype with certainty, or can predict the exact support a person needs. ADHD presentations are diverse. A test result without developmental and functional context can create more confidence than the information supports.

Why a comprehensive process can feel slower

A detailed assessment can seem less straightforward than a test with immediate graphs. It involves gathering records, completing questionnaires, speaking with family members where appropriate and allowing time to consider overlapping issues. Yet this process is designed to avoid reducing complex experiences to a single measurement.

For parents, this may mean the assessment explores more than classroom concentration or activity level. It may include sleep routines, learning progress, sensory experiences, mood, behaviour at home and school expectations. For adults, it may include periods when functioning was easier as well as periods when demands became harder to manage. These are not distractions from the assessment. They help build an accurate account of the person’s development and current needs.

A diagnosis is not the only useful outcome. Sometimes assessment identifies another explanation for the reported difficulties, or a combination of factors that requires support. Sometimes it confirms ADHD-related needs while also clarifying learning, anxiety, sleep or family factors that should be included in planning.

What happens after assessment

When ADHD is diagnosed or diagnostic support is provided, the next step is usually a multimodal plan. This may include psychological therapy, behavioural strategies,

support for executive functioning, school or workplace adjustments, family education and coordination with a GP, paediatrician or psychiatrist where medical decisions are needed.

Registered psychologists can provide assessment, therapy and psychological support. They do not prescribe medication. A prescriber may consider medical options as part of broader care, based on their own assessment and clinical responsibilities.

At Behavioural Neurotherapy Clinic, ADHD assessment is based on developmental history, structured measures, clinical interview and functional information rather than QEEG as a diagnostic test. Appointments are available in Doncaster for people across metropolitan Melbourne, with Australia-wide telehealth available where appropriate.

The useful question is not whether a brain map can provide a fast answer. It is whether the assessment creates a clear, evidence-informed pathway for understanding the difficulties and planning support that fits the person's everyday life.

For more information

Call (03) 9848 9100

Published: 14/11/2026 and updated on: 14/11/2026 by Dr Jacques Duff-PhD; BA Psych; Grad Dip Applied Psychology; FANSA; Clinical Neuroscientist