What Multimodal ADHD Planning Can Include



A diagnosis can explain a long history of missed deadlines, unfinished tasks, mental overload or difficulty regulating emotions. It does not, by itself, set out what happens on Monday morning at work, at home or when several demands arrive at once. Multimodal ADHD planning is the process of turning assessment findings into a practical, ongoing pathway that considers the person's daily environment, goals, mental health and medical care.

For adults, this planning often matters as much as the diagnostic process. Many have spent years using considerable effort to manage demands that appear straightforward to others. A useful plan does not assume that effort is the missing ingredient. It identifies where demands and available supports are out of balance, then sets out realistic ways to respond.

Multimodal ADHD planning is more than one appointment

Multimodal means using more than one form of support where it is relevant. It does not mean every person needs every option. The plan should be based on the assessment, current difficulties, personal priorities and the other professionals involved in care.

For one adult, the immediate priority may be creating workable systems for a role with competing deadlines. For another, sleep disruption, anxiety, low mood or relationship strain may need attention alongside executive-function difficulties. Someone returning to study may need support to understand access arrangements and organise tasks across a semester.

A psychologist’s role may include psychological assessment, therapy, behavioural strategies and coordination with relevant health professionals. A GP, paediatrician or psychiatrist manages medical decisions, including whether medication is appropriate. These roles are complementary. Clear communication helps avoid a plan in which each part of care operates separately.

The aim is not to make an adult with ADHD function as though they do not have ADHD. It is to build a support plan that accounts for the way attention, organisation, time awareness and emotional regulation can affect everyday life.

Start with the findings, not a generic checklist

A diagnosis is made through a process that considers more than a short list of current concerns. Adult ADHD assessment may include a developmental history, structured measures, information about education and employment, mental health

history, current functioning and, where available and appropriate, collateral information from someone who knew the person earlier in life.

This matters for planning. Difficulties with concentration can also occur alongside anxiety, depression, sleep problems, trauma-related symptoms, substance use, chronic stress or the effects of a concussion. Some adults have autistic traits, learning difficulties or other neurodevelopmental differences that also shape the support required. A plan needs to reflect that full picture rather than attribute every difficulty to ADHD.

Assessment findings can help identify patterns such as difficulty getting started, losing track of time, shifting between tasks too quickly, becoming stuck on a task, forgetting routine steps, or becoming overwhelmed when plans change. These are not character flaws. They are practical areas for discussion in therapy and daily planning.

Goals need to be specific enough to use

Broad goals such as being more organised can be hard to act on. Planning becomes more useful when the goal describes a real situation. This could mean arriving at work prepared for a morning handover, completing invoices before the end of the week, sharing household tasks more predictably, or starting an assignment without a late-night rush.

The goal also needs to fit the person’s circumstances. A complex colour-coded system may suit someone who enjoys detailed planning and has time to maintain it. It may be unhelpful for someone who already abandons systems once they require frequent updating. The most workable approach is often the one with the fewest steps between noticing a task and beginning it.

Psychological therapy and practical skill-building

Psychological therapy can provide a structured space to examine the thoughts, behaviours and situations that maintain day-to-day difficulties. Cognitive behavioural approaches may be used to work on patterns such as avoidance, all-or-nothing

thinking after a missed task, or the cycle of pressure and last-minute completion. Therapy may also address anxiety, low mood, sleep difficulties or emotional distress where these are part of the person's presentation.

Practical strategies are usually discussed alongside this work. Rather than relying on memory, a person may use one calendar, visual prompts, written task breakdowns or scheduled check-ins. Rather than setting aside an undefined block to do paperwork, the task may be narrowed to opening the document, finding the required information and completing the first section.

This is not about filling every hour with rules. Over-planning can become another barrier, particularly when a plan allows no room for travel, interruptions, fatigue or competing responsibilities. A clinician can help a person review which strategies are realistic, which create more pressure, and which need to be simplified.

Emotional regulation can also be part of the work. Some adults describe strong frustration, shame after mistakes, or difficulty recovering when plans change. Therapy may explore the triggers, early signs and responses that are relevant in that person's life. Where family members or partners are involved, sessions may also focus on communication, shared expectations and the practical division of responsibilities.

The environment belongs in the plan

ADHD support is not confined to an appointment room. Work, study, home routines and technology can either add demands or reduce them. Multimodal planning considers the setting in which a task must actually happen.

At work, helpful adjustments may involve written follow-up after verbal instructions, clearer priorities, quieter spaces for concentrated tasks, or breaking a large project into agreed milestones. The appropriate approach depends on the role, workplace policies and what the person wishes to disclose. Disclosure is a personal decision, not a requirement for planning.

At home, the focus may be on visible routines rather than intentions held in mind. A shared calendar, a consistent place for essential items, a weekly household meeting

or external reminders may reduce the number of decisions required at busy times. These strategies should be chosen with the household, rather than imposed on it.

For study, planning may include mapping assessment dates early, identifying the first action for each task, and discussing available support services with the education provider. When learning difficulties are also identified, the assessment findings may guide a more tailored discussion about study demands and adjustments.

Coordination with prescribers and other supports

Some adults choose to discuss medication with a GP or psychiatrist. BNC psychologists do not prescribe or make medical decisions. Where a prescriber is involved, psychological support can sit alongside medical care, with consent-based communication about the assessment findings, current goals and relevant changes in functioning.

Coordination may also involve a GP, psychiatrist, occupational therapist, dietitian, sleep clinician, school or workplace support person, depending on the individual situation. Not all of these are needed. The purpose is to make roles clear and reduce conflicting advice.

Nutritional assessment may be considered where indicated. This is different from using diet as a substitute for a full clinical plan. Changes to eating patterns, sleep routines or physical activity can be discussed in the context of the person’s health history, capacity and medical advice.

Review keeps the plan useful

A multimodal plan should be reviewed, particularly when circumstances change. A new job, a move, university exams, parenting demands, relationship stress or a period of poor sleep can expose gaps in a strategy that previously seemed manageable.

Review is not a test of whether someone has followed the plan perfectly. It is a chance to ask whether the plan fits the current demands. Sometimes the answer is to

add support. At other times, the better decision is to remove steps, lower the administrative load or focus on one priority before adding another.

For adults seeking support after an ADHD diagnosis, the next step does not need to be a complete overhaul of work, home and relationships. A structured plan can begin with the part of daily life creating the most pressure, while leaving room to adjust as circumstances become clearer. At BNC, ADHD therapy and multimodal planning can be discussed as part of an ongoing psychological support pathway.

For more information

Call (03) 9848 9100

Published: 17/11/2026 and updated on: 17/11/2026 by Dr Jacques Duff-PhD; BA Psych; Grad Dip Applied Psychology; FANSA; Clinical Neuroscientist